


AD-A056 338

STATE UNIV OF NEW YORK AT BUFFALO AMHERST STATISTICA--ETC F/8 12/1
THE INFORMATION MATRICES OF THE PARAMETERS OF MULTIPLE MIXED TI--ETC(U)
JUN 77 H J NEWTON N00014-75-C-0734
TR-53

UNCLASSIFIED

| OF |
AD
A056338

END
DATE
FILMED
8 -78
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

State University of New York at Buffalo
Department of Computer Science

STAT 100

4230 Ridge Lea Road, Room A58, Amherst, New York 14226

Telephone: 716-831-1225

LEVEL II

12

THE INFORMATION MATRICES OF THE PARAMETERS OF
MULTIPLE MIXED TIME SERIES*

by


H. Joseph Newton
State University of New York at Buffalo

AD A 056338

AD No. 1
DDC FILE COPY

TECHNICAL REPORT NO. 53

June 1977



DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

* Research supported in part by the Office of Naval Research.

78 07 12 012

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 53	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Information Matrices of the Parameters of Multiple Mixed Times Series		5. TYPE OF REPORT & PERIOD COVERED Technical rept.
7. AUTHOR(s) H. Joseph Newton		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Statistical Science Division State University of New York at Buffalo Amherst, New York 14226		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0734
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics and Probability Program Code 436 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR -042-234
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1977
		13. NUMBER OF PAGES 19 4222 P.
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/CONTINUING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multiple Time Series Information Matrices Autoregressive Schemes Moving Average Schemes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Closed form matrix equations are given for the information matrix of the parameters of the vector mixed autoregressive moving average time series model.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6501

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

449 511

78 07 12 012

THE INFORMATION MATRICES OF THE PARAMETERS OF
MULTIPLE MIXED TIME SERIES *

by

H. Joseph Newton
State University of New York at Buffalo
Amherst, New York

ABSTRACT

Closed form matrix equations are given for the information matrix of the parameters of the vector mixed autoregressive moving average time series model.

ACCESSION for	
NYIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION.....	
BY.....	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	APRIL 2nd or SPECIAL
A	

* Research supported in part by the Office of Naval Research.

1. Introduction

Consider the d-dimensional mixed autoregressive moving average process $\{\tilde{X}(t), t \in Z\}$, Z the set of integers, of order (p, q) ,

$$\sum_{j=0}^p A(j) \tilde{X}(t-j) = \sum_{k=0}^q B(k) \tilde{\epsilon}(t-k), \quad t \in Z$$

where $A(0) \equiv B(0) \equiv I_d$, the d-dimensional identity matrix, and $\{\tilde{\epsilon}(t), t \in Z\}$ is a collection of uncorrelated, zero mean, d-dimensional random variables each having positive definite covariance matrix Φ .

The estimation of the $(d \times d)$ matrices $A(1), \dots, A(p)$, $B(1), \dots, B(q)$, and Φ has received considerable attention in the recent literature (Akaike (1973a), Wilson (1973), Dunsmuir and Hannan (1976), for example). One of the most difficult computational problems involved in the estimation is that of determining the asymptotic covariance matrix V of the maximum likelihood estimators. Hannan (1970), p. 385 and p. 329 has given expressions for V for the case $p = 0$ or $q = 0$.

In this paper we derive closed form matrix expressions for V by using Whittle's (1953) formula for the Fisher information matrix of the parameters of a Gaussian multiple time series.

2. The Information Matrix of the Mixed Process

Let $\{\tilde{Y}(t), t \in \mathbb{Z}\}$ be a zero mean Gaussian time series whose distribution depends on parameters $\tilde{\theta} = (\theta_1, \dots, \theta_r)^T$, where A^T denotes the transpose of the matrix A . Let $f(\omega)$, $\omega \in [-\pi, \pi]$ be the spectral density matrix of $\tilde{Y}(\cdot)$. Then given a sample realization $y(1), \dots, y(T)$ of $\tilde{Y}(\cdot)$, Whittle (1953) shows that the maximum likelihood estimators $\hat{\tilde{\theta}}_T$ of $\tilde{\theta}$ are such that $\sqrt{T}(\hat{\tilde{\theta}}_T - \tilde{\theta})$ is asymptotically r -dimensional normal with mean zero and covariance matrix $V(\tilde{\theta}) = I^{-1}(\tilde{\theta})$ where the (j,k) th element of $I(\tilde{\theta})$ is given by

$$I_{jk}(\tilde{\theta}) = \frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr} \left[\frac{\partial f(\omega)}{\partial \theta_j} f^{-1}(\omega) \frac{\partial f(\omega)}{\partial \theta_k} f^{-1}(\omega) \right] d\omega, \quad (2.1)$$

where the notation $\frac{\partial A}{\partial b}$, b a scalar and $A = (A_{jk})$ and $(n \times m)$ matrix, denotes the $(n \times m)$ matrix of derivatives $\left(\frac{\partial A_{jk}}{\partial b} \right)$, and $\text{tr} B$ denotes the trace of the matrix B .

Let $D = (D_{jk})$ be an $(n \times n)$ matrix and $U = U(D)$, $V = V(D)$ be matrix functions of D . Then (Neudecker (1969), for example)

$$\frac{\partial D}{\partial D_{jk}} = E_{jk}, \quad \frac{\partial D^T}{\partial D_{jk}} = E_{jk}^T = E_{kj}, \quad (2.2)$$

where E_{jk} is the zero matrix except the (j,k) th element which is one.

Also, the chain rule holds;

$$\frac{\partial UV}{\partial D_{jk}} = \frac{\partial U}{\partial D_{jk}} V + U \frac{\partial V}{\partial D_{jk}}. \quad (2.3)$$

Thus from $D^{-1}D = I_n$, where I_n denotes the n th order identity matrix, we have

$$\frac{\partial D^{-1}}{\partial a} = -D^{-1} \frac{\partial D}{\partial a} D^{-1} \quad (2.4)$$

Note that the trace operation satisfies

$$\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$$

$$\text{tr}(AB) = \text{tr}(BA)$$

From (2.4) we can write (2.1) as

$$I_{jk} = \frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr} \left[\frac{\partial f^{-1}}{\partial \theta_j} f \frac{\partial f^{-1}}{\partial \theta_k} f \right] d\omega \quad (2.5)$$

$$= -\frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr} \left[\frac{\partial f^{-1}}{\partial \theta_j} \frac{\partial f}{\partial \theta_k} \right] d\omega, \quad (2.6)$$

where we have now deleted the argument of $f(\cdot)$ for convenience.

The spectral density matrix $f(\cdot)$ of the mixed model can be written (see Hannan (1970), p. 67)

$$f(\omega) = \frac{1}{2\pi} G^{-1}(e^{i\omega}) H(e^{i\omega}) \dagger H^*(e^{i\omega}) G^{-*}(e^{i\omega}), \quad \omega \in [-\pi, \pi] \quad (2.7)$$

where the complex matrix polynomials $G(\cdot)$ and $H(\cdot)$ are given by

$$G(z) = \sum_{j=0}^P A(j) z^j$$

$$H(z) = \sum_{k=0}^q B(k) z^k ,$$

and A^* denotes the complex conjugate transpose of the matrix A . We assume that the zeros of $\det(G(z))$ and $\det(H(z))$ are outside the unit circle so that the elements of $G^{-1}(z)$ and $H^{-1}(z)$ can be written as power series in z .

From (2.7) we write

$$2\pi f = G^{-1} H \dagger H^* G^{-*} \equiv G^{-1} Q G^{-*} \quad (2.8)$$

$$\frac{1}{2\pi} f^{-1} = G^* Q^{-1} G , \quad (2.9)$$

where G doesn't involve the $B(\cdot)$, $Q = H \dagger H^*$ is Hermitian and mathematically independent of the $A(\cdot)$, $Q = \dagger$ if $q = 0$, and we have deleted the arguments of all functions.

Clearly we must order the elements of the $A(\cdot)$ and $B(\cdot)$ into a vector $\underline{\theta}$. However, we first find the element of the information matrix corresponding to $A_{jk}(v)$, $A_{lm}(u)$ (denoted $I[A_{jk}(v), A_{lm}(u)]$) by (2.5), the element corresponding to $B_{jk}(v)$, $B_{lm}(u)$ (denoted $I[B_{jk}(v), B_{lm}(u)]$) by (2.1), and the element corresponding to $A_{jk}(v)$, $B_{lm}(u)$ (denoted $I[A_{jk}(v), B_{lm}(u)]$ by (2.6)). Then we consider various orderings of the elements of the $A(\cdot)$ and $B(\cdot)$ to find a convenient expression for $I(\underline{\theta})$.

$$\underline{I[A_{jk}(v), A_{lm}(u)]}$$

From (2.2), (2.3), and (2.5) we obtain

$$\begin{aligned} \frac{1}{2\pi} \frac{\partial f^{-1}}{\partial A_{jk}(v)} &= \frac{\partial}{\partial A_{jk}(v)} G^* Q^{-1} G \\ &= \frac{\partial G^* Q^{-1}}{\partial A_{jk}(v)} G + G^* Q^{-1} \frac{\partial G}{\partial A_{jk}(v)} \\ &= E_{kj} Q^{-1} G e^{-iv\omega} + G^* Q^{-1} E_{jk} e^{iv\omega} . \end{aligned}$$

Thus

$$\frac{\partial f^{-1}}{\partial A_{jk}(v)} f = E_{kj} G^{-*} e^{-iv\omega} + f^{-1} G^{-1} E_{jk} f e^{iv\omega} ,$$

and

$$\begin{aligned} \frac{\partial f^{-1}}{\partial A_{jk}(v)} f \frac{\partial f^{-1}}{\partial A_{lm}(u)} f &= E_{kj} G^{-*} E_{ml} G^{-*} e^{-i(u+v)\omega} + 2\pi E_{kj} Q^{-1} E_{lm} f e^{-i(v-u)\omega} \\ &\quad + 2\pi G^* Q^{-1} E_{jk} f E_{ml} G^{-*} e^{-i(u-v)\omega} \\ &\quad + f^{-1} G^{-1} E_{jk} G^{-1} E_{lm} f e^{i(u+v)\omega} . \end{aligned}$$

Denote by D^{rs} the matrix D replaced by zeros except for the r^{th} row which is replaced by the s^{th} row, i.e. $D^{rs} = E_{rs} D$. Then

for matrices $D = (D_{jk})$, $C = (C_{lm})$ we have

$$\text{tr}[E_{rs}^D E_{tu}^C] = \text{tr}[D^{rs} C^{tu}] = D_{st} C_{ur}$$

Thus

$$\begin{aligned} \text{tr} \left[\frac{\partial f^{-1}}{\partial A_{jk}(v)} f \frac{\partial f^{-1}}{\partial A_{lm}(u)} f \right] &= \text{tr}[E_{kj} G^{-*} E_{ml} G^{-*}] e^{-i(u+v)\omega} \\ &+ 2\pi \text{tr}[E_{kj} Q^{-1} E_{lm} f] e^{-i(v-u)\omega} \\ &+ 2\pi \text{tr}[E_{ml} Q^{-1} E_{jk} f] e^{-i(u-v)\omega} \\ &+ \text{tr}[E_{lm} G^{-1} E_{jk} G^{-1}] e^{i(u+v)\omega} \\ &= G_{jm}^{-*} G_{lk}^{-*} e^{-i(u+v)\omega} + 2\pi Q_{jl}^{-1} f_{mk} e^{-i(v-u)\omega} \\ &+ 2\pi Q_{lj}^{-1} f_{km} e^{-i(u-v)\omega} + G_{mj}^{-1} G_{kl}^{-1} e^{i(u+v)\omega} . \end{aligned}$$

We argue that the first and last terms integrate to zero, as follows:

because the roots of $\det(G(z)) = 0$ are assumed strictly outside the unit circle, G_{jk}^{-1} , the (j,k) th element of G^{-1} , can be written as a power series (which allows the interchange of summation and integration in (2.10) below)

$$G_{jk}^{-1}(z) = \sum_{\ell=0}^{\infty} C_{jk}(\ell) z^{\ell}$$

and thus

$$\begin{aligned} & \int_{-\pi}^{\pi} G_{mj}^{-1} G_{k\ell}^{-1} e^{i(u+v)\omega} d\omega \\ &= \int_{-\pi}^{\pi} \sum_{t=0}^{\infty} \sum_{r=0}^{\infty} C_{mj}(t) C_{k\ell}(r) e^{i(u+v+t+r)\omega} d\omega \quad (2.10) \\ &= \sum_{t=0}^{\infty} \sum_{r=0}^{\infty} C_{mj}(t) C_{k\ell}(r) \int_{-\pi}^{\pi} e^{i(u+v+t+r)\omega} d\omega, \end{aligned}$$

and the integral is always zero since $(u+v+t+r) > 0$.

Thus we are left with

$$I[A_{jk}(v), A_{\ell m}(u)] = \frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr}[C + C^*] d\omega,$$

where

$$C = 2\pi E_{kj} Q^{-1} E_{\ell m} f e^{-i(v-u)\omega}.$$

Since the integral is real, we have

$$\begin{aligned} I[A_{jk}(v), A_{\ell m}(u)] &= \frac{1}{4\pi} \int_{-\pi}^{\pi} 2 \text{tr}[C] d\omega \\ &= \int_{-\pi}^{\pi} Q_{jl}^{-1} f_{mk} e^{-i(v-u)\omega} d\omega. \end{aligned} \quad (2.11)$$

$$\underline{I[B_{jk}(v), B_{lm}(u)]}$$

From (2.8) we have

$$\begin{aligned} 2\pi \frac{\partial f}{\partial B_{jk}(v)} &= \frac{\partial}{\partial B_{jk}(v)} G^{-1} H \nabla H^* G^{-*} \\ &= \frac{\partial G^{-1} H}{\partial B_{jk}(v)} \nabla H^* G^{-*} + G^{-1} H \frac{\partial \nabla H^* G^{-*}}{\partial B_{jk}(v)} \\ &= G^{-1} E_{jk} \nabla H^* G^{-*} e^{i v \omega} + G^{-1} H \nabla E_{kj} G^{-*} e^{i v \omega} . \end{aligned}$$

Then

$$\begin{aligned} \frac{\partial f}{\partial B_{jk}(v)} f^{-1} \frac{\partial f}{\partial B_{lm}(u)} f^{-1} &= G^{-1} E_{jk} \nabla H^* Q^{-1} E_{lm} \nabla H^* Q^{-1} G e^{i(u+v)\omega} \\ &+ G^{-1} E_{jk} \nabla H^* Q^{-1} H \nabla E_{ml} Q^{-1} G e^{-i(u-v)\omega} \\ &+ G^{-1} H \nabla E_{kj} Q^{-1} E_{lm} \nabla H^* Q^{-1} G e^{-i(v-u)\omega} \\ &+ G^{-1} H \nabla E_{kj} Q^{-1} H \nabla E_{ml} Q^{-1} G e^{-i(u+v)\omega} . \end{aligned}$$

The trace of this is given by

$$\text{tr} \left[\frac{\partial f}{\partial B_{jk}(v)} f^{-1} \frac{\partial f}{\partial B_{lm}(u)} f^{-1} \right] = \text{tr}[C + C^*] + \text{tr}[D + D^*] ,$$

where

$$C = E_{\ell m} H^{-1} E_{jk} H^{-1} e^{i(u+v)\omega}$$

$$D = E_{m\ell} Q^{-1} E_{jk} \int e^{-i(u-v)\omega}$$

The integral of the first term again vanishes, and we have

$$I[B_{jk}(v), B_{\ell m}(u)] = \frac{1}{2\pi} \int_{-\pi}^{\pi} Q_{j\ell}^{-1} \int_{\ell}^{\pi} e^{-i(v-u)\omega} d\omega \quad (2.12)$$

$$\underline{I[A_{jk}(v), B_{\ell m}(u)]}$$

To find the off block diagonal elements of I we obtain, using arguments similar to the ones above, that

$$\text{tr} \left[\frac{\partial f^{-1}}{\partial A_{jk}(v)} \frac{\partial f}{\partial B_{\ell m}(u)} \right] = \text{tr}[C + C^*] + \text{tr}[D + D^*] ,$$

where

$$C = E_{\ell m} H^{-1} E_{jk} G^{-1} e^{i(u+v)\omega}$$

$$D = E_{kj} Q^{-1} E_{\ell m} \int H^* G^{-*} e^{-i(v-u)\omega} .$$

Thus

$$I[A_{jk}(v), B_{\ell m}(u)] = \frac{1}{2\pi} \int_{-\pi}^{\pi} Q_{j\ell}^{-1} \left[\int_{\ell}^{\pi} H^* G^{-*} \right]_{mk} e^{-i(v-u)\omega} d\omega , \quad (2.13)$$

where the (m,k) th element of the product $\ddagger H^* G^{-*}$ is denoted $[\ddagger H^* G^{-*}]_{mk}$.

So the integrands in the expressions for $I[A_{jk}(v), A_{lm}(u)]$, $I[B_{jk}(v), B_{lm}(u)]$, and $I[A_{jk}(v), B_{lm}(u)]$ are all of the form $C_{jl}^{-1} D_{mk} e^{-i(v-u)\omega}$ for some matrices C and D , i.e., we have the transformation of indices denoted

$$(j,k,v), (\ell,m,u) \rightarrow (j,\ell,-v), (m,k,u) \quad (2.14)$$

This transformation allows us to find general expressions for the information matrix for various orderings of the elements of the $A(\cdot)$ and $B(\cdot)$ matrices.

Ordering of $A(\cdot)$, $B(\cdot)$

Define the partitioned matrices

$$A_1 = \begin{pmatrix} A(1) & \vdots & \dots & \vdots & A(p) \end{pmatrix},$$

$$A_2 = \begin{pmatrix} A^T(1) & \vdots & \dots & \vdots & A^T(p) \end{pmatrix},$$

$$B_1 = \begin{pmatrix} B(1) & \vdots & \dots & \vdots & B(q) \end{pmatrix},$$

$$B_2 = \begin{pmatrix} B^T(1) & \vdots & \dots & \vdots & B^T(q) \end{pmatrix}.$$

Then let

$$\alpha_1 = \text{vec} (A_1)$$

$$\alpha_2 = \text{vec} (A_1^T)$$

$$\alpha_3 = \text{vec} (A_2)$$

$$\beta_1 = \text{vec} (B_1^T)$$

$$\beta_2 = \text{vec} (B_2^T)$$

$$\beta_3 = \text{vec} (B_1)$$

$$J_{nm}(\omega) = (J_{jk}(\omega)) = e^{i(j-k)\omega}, \quad j = 1, \dots, n, k = 1, \dots, m,$$

where $\text{vec} (A)$ denotes the operation of stacking the columns of A , i.e., if $A = (a_1, \dots, a_n)$, $\text{vec} (A) = (a_1^T \vdots \dots \vdots a_n^T)^T$.

Define the Kronecker product of the $n \times m$ matrix $A = (A_{jk})$ and the $r \times s$ matrix $B = (B_{jk})$ as the $nr \times ms$ matrix

$$C = A \otimes B = \begin{bmatrix} A_{11}B & \dots & A_{1m}B \\ \vdots & & \vdots \\ A_{n1}B & \dots & A_{nm}B \end{bmatrix}.$$

Denote the information matrix of parameters θ_1, θ_2 by $I^{\theta_1 \theta_2}$. Note that $I^{\theta_2 \theta_1} = (I^{\theta_1 \theta_2})^T$, and that $(C \otimes D)^T = C^T \otimes D^T$. Then we have from (2.11), (2.12), and (2.13),

$$I_{\alpha_1 \alpha_1} = \int_{-\pi}^{\pi} J_{pp}^T \otimes f^T \otimes Q^{-1} d\omega$$

$$I_{\alpha_2 \alpha_2} = \int_{-\pi}^{\pi} Q^{-T} \otimes J_{pp} \otimes f d\omega$$

$$I_{\alpha_3 \alpha_3} = \int_{-\pi}^{\pi} J_{pp} \otimes Q^{-T} \otimes f d\omega$$

$$I_{\beta_1 \beta_1} = \frac{1}{2\pi} \int_{-\pi}^{\pi} J_{qq} \otimes Q^{-T} \otimes \ddagger d\omega$$

$$I_{\beta_2 \beta_2} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \ddagger \otimes J_{qq} \otimes Q^{-T} d\omega$$

$$I_{\beta_3 \beta_3} = \frac{1}{2\pi} \int_{-\pi}^{\pi} J_{qq}^T \otimes \ddagger \otimes Q^{-1} d\omega$$

$$I_{\alpha_1 \beta_3} = \frac{1}{2\pi} \int_{-\pi}^{\pi} J_{qp}^T \otimes (\ddagger H^* G^{-*})^T \otimes Q^{-1} d\omega$$

We have derived the information matrices for three methods of ordering the elements of the parameter matrices. The results for α_1 and β_3 are used below to find a convenient form for the information matrix of the parameters of a mixed scheme. The results for α_1 and α_2 are used for autoregressive processes, while those for β_1 are used for moving average processes. To summarize, we have

Remarks on Information Matrices

$$\begin{aligned} \text{a) } I &= \begin{bmatrix} I_{\alpha_1 \alpha_1} & I_{\alpha_1 \beta_3} \\ \text{sym } I & I_{\beta_3 \beta_3} \end{bmatrix} = \int_{-\pi}^{\pi} \begin{bmatrix} J_{pp}^T \otimes f^T \otimes Q^{-1} & -\frac{1}{2\pi} J_{qp}^T \otimes (\ddagger H^* G^{-*})^T \otimes Q^{-1} \\ \text{sym} & \frac{1}{2\pi} J_{qq}^T \otimes \ddagger \otimes Q^{-1} \end{bmatrix} d\omega \\ &= \int_{-\pi}^{\pi} \begin{pmatrix} X \\ Y \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}^* d\omega, \end{aligned}$$

where

$$X = (2\pi)^{-\frac{1}{2}} J_{\sim p} \otimes (\Sigma^{\frac{1}{2}} H^* G^{-*})^T \otimes H^{-*} \Sigma^{-\frac{1}{2}}$$

$$Y = -(2\pi)^{-\frac{1}{2}} J_{\sim q} \otimes \Sigma^{\frac{1}{2}} \otimes H^{-*} \Sigma^{-\frac{1}{2}},$$

$\Sigma^{\frac{1}{2}}$ denotes a positive definite square root of the positive definite matrix Σ , and $J_{\sim r}$ is an r -dimensional vector whose k th element is $e^{-ik\omega}$.

b) To find the information matrix of the elements of Σ , and the cross information, in turn, between the $A(\cdot)$ and Σ , and the $B(\cdot)$ and Σ , we note that since Σ is symmetric

$$\frac{\partial f}{\partial \Sigma_{jk}} = \frac{1}{2\pi} C[E_{jk} + (1 - \delta_{jk}) E_{kj}] C^*,$$

where $C = G^{-1}H$, and $\delta_{jk} = 1$ if $j = k$ and 0 otherwise. Defining

$$D_{jk} = E_{jk} + (1 - \delta_{jk}) E_{kj},$$

we have

$$\text{tr} \left[\frac{\partial f}{\partial \Sigma_{jk}} f^{-1} \frac{\partial f}{\partial \Sigma_{lm}} f^{-1} \right] = \text{tr} [D_{jk} \Sigma^{-1} D_{lm} \Sigma^{-1}],$$

which is independent of ω . Thus the element of the information matrix corresponding to Σ_{jk}, Σ_{lm} , denoted $I[\Sigma_{jk}, \Sigma_{lm}]$, is given by

$$\begin{aligned}
I[\mathbb{X}_{jk}, \mathbb{X}_{lm}] &= \frac{1}{2} \text{tr}[D_{jk} \mathbb{X}^{-1}] \\
&= \frac{1}{2} \left[(2 - \delta_{jk} - \delta_{lm} + \delta_{jk} \delta_{lm}) \mathbb{X}_{jl}^{-1} \mathbb{X}_{mk}^{-1} \right. \\
&\quad \left. + (2 - \delta_{jk} - \delta_{lm}) \mathbb{X}_{kl}^{-1} \mathbb{X}_{mj}^{-1} \right], \quad (2.15)
\end{aligned}$$

where we consider $\mathbb{X}_{u,v}$, $u \geq v = 1, \dots, d$ as the distinct elements of \mathbb{X} . To write this in matrix form, define

$$\sigma_1 = \text{vec}(\mathbb{X})$$

$$\sigma_2 = \text{lvec}(\mathbb{X}),$$

where $\text{lvec}(A)$ is the vec operation on the lower triangular portion of the matrix A . Thus σ_2 contains the distinct elements of \mathbb{X} .

If we treat the d^2 elements of \mathbb{X} as being distinct (i.e., ignore the symmetry of \mathbb{X}), then $D_{jk} = E_{jk}$, and it is easy to show that

$$I^{\sigma_1 \sigma_1} = \frac{1}{2} \mathbb{X}^{-1} \otimes \mathbb{X}^{-1}.$$

Note that we can write

$$\sigma_1 = A \sigma_2$$

where A is a $d^2 \times \frac{d(d+1)}{2}$ matrix of zeros and ones. Then
(Pagano (1974))

$$\begin{aligned} I_{\sigma_2 \sigma_2} &= A^T I_{\sigma_1 \sigma_1} A \\ &= \frac{1}{2} A^T (\Phi^{-1} \otimes \Phi^{-1}) A \end{aligned} \quad (2.16)$$

Thus the information matrix of σ_1 can be found from either (2.15) or (2.16).

c) One can also show that

$$\begin{aligned} I[A_{jk}(v), \Phi_{lm}] &= \frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr} \left[\frac{\partial f}{\partial A_{jk}(v)} f^{-1} \frac{\partial f}{\partial \Phi_{lm}} f^{-1} \right] d\omega \\ &= 0, \end{aligned}$$

and

$$\begin{aligned} I[B_{jk}(v), \Phi_{lm}] &= \frac{1}{4\pi} \int_{-\pi}^{\pi} \text{tr} \left[\frac{\partial f}{\partial B_{jk}(v)} f^{-1} \frac{\partial f}{\partial \Phi_{lm}} f^{-1} \right] d\omega \\ &= 0. \end{aligned}$$

These integrals are zero for the same reason that the integrals vanish in the derivation of the information matrices of the $A(\cdot)$ and the $B(\cdot)$.

d) If $q = 0$ we have

$$I^{\alpha_1 \alpha_1} = \Gamma_p \otimes \mathbb{I}^{-1} \quad (2.17)$$

$$I^{\alpha_2 \alpha_2} = \mathbb{I}^{-1} \otimes \Gamma_p, \quad (2.18)$$

since

$$\Gamma_p \equiv \text{BTOEPL}(R(0), R(-1), \dots, R(1-p))$$

$$= \int_{-\pi}^{\pi} J_{pp} \otimes f \, d\omega,$$

where $\text{BTOEPL}(R(0), \dots, R(1-p))$ is a block Toeplitz matrix having

$R(j-k)$ in the j th row and k th column of blocks, $j, k = 1, \dots, p$.

Note that (2.18) is the result given by Hannan (1970), p. 329.

e) If $p = 0$, then $Q^{-T} = \frac{1}{2\pi} f^{-T}$ which is of the form of an autoregressive spectral density matrix. Thus $I^{\beta_1 \beta_1}$ should be of the same form as $I^{\alpha_1 \alpha_1}$ of the previous note with the autocovariances corresponding to the spectral density Q^{-T} replacing $R(\cdot)$ and $2\pi \mathbb{I}^{-1}$ replacing \mathbb{I} . Thus

$$I^{\beta_1 \beta_1} = \Gamma_{1q} \otimes \mathbb{I} \quad (2.19)$$

where

$$\Gamma_{\mathbf{i}_q} = \text{BTOEPL}(\text{Ri}(0), \dots, \text{Ri}(1-q)) ,$$

$$\text{Ri}(\mathbf{v}) = \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \mathbf{f}^{-T}(\omega) e^{i\mathbf{v}\omega} d\omega , \quad \mathbf{v} \in \mathbb{Z} .$$

See Newton (1975), p. 11, et seq for a discussion of the inverse autocovariances $\text{Ri}(\cdot)$. Note that (2.19) is the result stated by Hannan (1970), p. 385.

f) Equations (2.18) and (2.19) also provide an interpretation of information matrices of random matrices (transformed into a random vector by the vec operator) as the covariance matrix of some random vector. We can write

$$\mathbf{I}^{\alpha_1 \alpha_1} = \{\text{R}(\mathbf{j} - \mathbf{k}) \otimes \mathbf{I}_p^{-1}\}_{p \times p}$$

and

$$\mathbf{I}^{\beta_1 \beta_1} = \{\text{Ri}(\mathbf{j} - \mathbf{k}) \otimes \mathbf{I}_q^{-1}\}_{q \times q} ,$$

i.e., $\mathbf{I}^{\alpha_1 \alpha_1}$ is a $(p \times p)$ block matrix of $(d^2 \times d^2)$ blocks, while $\mathbf{I}^{\beta_1 \beta_1}$ is a $(q \times q)$ block matrix of $(d^2 \times d^2)$ blocks. Thus there exist d^2 -dimensional random vectors $\mathbf{z}_1, \dots, \mathbf{z}_p$ and $\mathbf{y}_1, \dots, \mathbf{y}_q$ such that $\mathbf{z} = \begin{pmatrix} \mathbf{z}_1^T : \dots : \mathbf{z}_p^T \end{pmatrix}^T$ had covariance matrix given by $\mathbf{I}^{\alpha_1 \alpha_1}$ and $\mathbf{y} = \begin{pmatrix} \mathbf{y}_1^T : \dots : \mathbf{y}_q^T \end{pmatrix}^T$ has covariance matrix given by $\mathbf{I}^{\beta_1 \beta_1}$.

g) Akaike (1973a) derives approximate expressions for the elements of the Hessian of the log likelihood function. From the formulas (2.11),

(2.12), (2.13) above, it is clear that Akaike's formulas for the Hessian are the sample analogues of the information matrix of the mixed scheme parameters. Thus one can use the block Toeplitz matrix inversion techniques developed by Akaike (1973b) to find the asymptotic covariance matrix of the maximum likelihood estimators.

Acknowledgments

The author would like to thank Professors E. Parzen and M. Pagano for many helpful discussions.

References

- Akaike, H. (1973a), "Maximum likelihood identification of Gaussian autoregressive moving average models," Biometrika, 60, 255-265.
- Akaike, H. (1973b), "Block Toeplitz matrix inversion," SIAM J. App. Math., 24, 234-241.
- Dunsmuir, W. and Hannan, E. J. (1976), "Vector linear time series models," Advancements in Applied Probability, 8, 339-364.
- Hannan, E. J., (1970), Multiple Time Series, New York, John Wiley.
- Neudecker, H. (1969), "Some theorems on Matrix differentiation with special reference to Kronecker matrix products," Journal of the American Statistical Association, 64, 953-964.
- Newton, H. J. (1975), "The efficient estimation of stationary multiple time series mixed models: theory and algorithms," Technical Report No. 33, Statistical Science Division, SUNY at Buffalo.
- Pagano, M. (1974), "Modification of parameter estimates under linear constraints," Technical Report No. 12, Statistical Science Division, SUNY at Buffalo.
- Rao, C. R. (1973), Linear Statistical Inference and Its Applications, (2nd ed), New York, John Wiley.
- Whittle, P. (1953), "The analysis of multiple stationary time series," Journal of the Royal Statistical Society B, 15, 125-139.
- Wilson, G. T. (1973), "The estimation of parameters in multivariate time series models," Journal of the Royal Statistical Society B, 35, 76-85.